

# UV Complete QFTs by $[T \bar{T}]_s$ deformations

**Changrim Ahn**

Ewha Womans University



**Mathematics and Physics of Integrability (MPI2024)**

Matrix Institute, U of Melbourne

## Goal

Complete Classifications of UV Complete 2D QFTs

which RG flow into a given IR CFT

### PART 1. RG flows from IR to UV

[with Andre LeClair (2022)]

- Massless  $S$ -matrices of CFTs deformed by  $(T\bar{T})_{s+1}$
  - Complete classifications based on “Plateaux equations” (TBA in UV limit)
- 

### PART 2. Identifying UV complete QFTs

[with Zoltan Bajnok (2024)]


- Apply to minimal CFTs  $\mathcal{M}_p$
- Identify UV QFTs from Reflection amplitudes vs. TBA

# PART 1. RG flows from IR to UV

## Conventional Approach (top down)

- UV CFT + a relevant field  $\rightarrow$  IR CFT + irrelevant fields
  - Only a special relevant field maintains the integrability
  - Common in CMP since physics at low T is important to find non-trivial (Wilson-Fisher) fixed points
- (Ex) Zamolodchikov Flows originally based on conjectured TBA

$$\mathcal{M}_{p+1} + \lambda \Phi_{1,3} \rightarrow \mathcal{M}_p + \lambda' \Phi_{3,1}, \quad p = 3, 4, \dots$$



- Many more RG flows [Dorey, Dunning, Ravanini, ...]
  - have been conjectured based on TBA or NLIE
  - Lagrangians / exact S-matrices are missing

## Our new approach (bottom up) [CA, A.LeClair (2022)]

- IR CFT + irrelevant fields  $\mapsto$  UV CFT + a relevant field
- Natural since  $S$ -matrices are defined in the IR (infinite volume)
- Common in HEP where UV complete theory is being searched (GUT, SUSY, Superstring, ...)
- Use  $T\bar{T}$  as a ladder

## $T\bar{T}$ deformations [Tateo's talk]

- Very active developments [Tateo, Zamolodchikov, ...]
- Energy-momentum tensor  $T_2$
- All higher conserved charges =  $\{ [T\bar{T}]_s \}$

$$[T\bar{T}]_s = T_{s+1}\bar{T}_{s+1} - \Theta_{s-1}\bar{\Theta}_{s-1}, \quad \partial_{\bar{z}}T_{s+1} = \partial_z\Theta_{s-1}$$

- Preserve integrability
- Exact results possible for even non-integrable theories

## Space of 2D IQFTs [Smirnov-Zamolodchikov (2017)]

- Expands the integrable space in infinite dimensions
- If the mother theory is integrable, new integrable QFTs

$$\text{New IQFTs} = \text{an IQFT} + \sum_{s=1}^{\infty} \alpha_s [T\bar{T}]_s$$

- Exact  $S$ -matrices are given by additional CDD factors

$$S(\theta) = \prod_{s=1}^{\infty} e^{2i\alpha_s M^{2s} \sinh(s\theta)} \cdot S_0(\theta)$$



## Swampland (Hagedorn singularity) to cross for UV

- Burgers Equation ( $s = 1$ ) [ $CFT + \alpha_1[T\bar{T}]_1$ ]

$$\partial_\alpha E + E \partial_R E = 0 \quad \rightarrow \quad c_{\text{eff}}(R) = \frac{2c_0}{1 + \sqrt{1 + \frac{2\pi\alpha_1 c_0}{3R^2}}}$$

with square root singularity at  $R_c = \sqrt{\frac{2\pi|\alpha_1|c_0}{3}}$

- Singularity occurs also for each  $[T\bar{T}]_s$  with  $s \geq 1$
- We show that the singularities can be avoided if we fine-tune all  $\alpha_s$

## S-matrices of CFTs from a massless limit

- Consider “massive” integrable deformation of a CFT e.g.

$$[G]_k \equiv \frac{G_1 \otimes G_k}{G_{k+1}} + \lambda \Phi_{\text{least rel}}$$

- Integrable with exact  $S$ -matrix  $S_0$  [Bernard, LeClair, CA (1990)]
- Take  $\lambda \rightarrow 0^-$  or  $M \rightarrow 0$  limit
  - rescale the rapidity

$$M \rightarrow 0 \ \& \ \theta = \pm\Lambda + \hat{\theta} \quad \text{with finite } Me^\Lambda = \mu$$

- $R$  and  $L$  particles appear depending on  $\pm$

$$(R) : E = P = \frac{\mu}{2} e^{\hat{\theta}}, \quad (L) : E = -P = \frac{\mu}{2} e^{-\hat{\theta}}$$

- $S$ -matrices between  $RR, LL$  are the same  $S^{RR}(\hat{\theta}) = S^{LL}(\hat{\theta}) = S_0(\hat{\theta})$
- $S$ -matrices between  $RL, LR$  are trivial since  $\theta_{12} \rightarrow \pm\infty$

## S-matrices for deformed CFTs by $T\bar{T}$ 's

- CDD factors become trivial in  $RR, LL$  sectors

$$\prod_{s=1}^{\infty} e^{2i\alpha_s M^{2s} \sinh(s\theta)} \rightarrow 1 \quad \text{as } M \rightarrow 0$$

- CDD factors become non-trivial in  $RL, LR$  sectors

$$\prod_{s=1}^{\infty} e^{2i\alpha_s M^{2s} \sinh(s\theta)} \rightarrow \prod_{s=1}^{\infty} e^{\pm i\alpha_s \mu^{2s} e^{\pm s\hat{\theta}}} \equiv S_{\text{CDD}}(\hat{\theta})$$

- Summary

$$S^{RR}(\theta) = S^{LL}(\theta) = S_0, \quad S^{RL}(\theta) = S^{LR}(-\theta) = S_{\text{CDD}}(\theta)$$

## How to restrict $S_{\text{CDD}}$ ?

- Find all possible  $S_{\text{CDD}}$  for a given  $S_0$  which satisfy
- UV completeness
  - No Hagedorn singularities
  - $c_{\text{eff}}$  in the UV limit is **finite, rational**
- Crossing-Unitarity

$$S_{\text{CDD}}(\theta)S_{\text{CDD}}(\theta + i\pi) = 1$$

from which  $S_{\text{CDD}}$  are arbitrary products of only two factors

$$S_{\text{CDD}}^{(1)} = -\tanh\left(\frac{\theta - \beta}{2} - \frac{i\pi}{4}\right), \quad S_{\text{CDD}}^{(2)} = \frac{\sinh\theta - i\sin\beta}{\sinh\theta + i\sin\beta}$$

- All  $\alpha_s$  are fixed exactly in terms of  $\beta$ 's
- We can work with UV limit of TBA, namely, “plateaux equations”

## Focus on “diagonal” $S$ -matrices theories

- Affine-Toda theories [Braden, Corrigan, Dorey, Sasaki (1990)] ( $k = 1$  cases of the coset CFTs  $[G]_1$ )
- Coupling constant independent part of ATFT (diagonal)  $S$ -matrices provide  $S_0^{ab}$
- A CFT can have different  $S$ -matrices depending on the choice of the integrable relevant field (ex) Ising CFT by  $G = su(2)$ ,  $E_8$
- Plateaux equations

$$x_a^R = \prod_{b=1}^r (1 + x_b^R)^{k_{ab}} \prod_{b=1}^r (1 + x_b^L)^{\tilde{k}_{ab}}$$

where exponents are integrals of logarithmic derivatives of  $S$ -matrices  $S_0^{ab}$  and  $S_{\text{CDD}}^{ab}$  ( $\tilde{k}_{ab} = \frac{1}{2}, 1$  or sum of them)

- Expressions in terms of universal kernel are not available
- $c_{\text{eff}}$  in terms of some dilog identities

## Results: UV CFTs which can flow into Ising CFT $\mathcal{M}_3$

- From the Plateaux equations, we find only

$$c_{\text{eff}} = \frac{7}{10}, \frac{21}{22}, \frac{3}{2}, \frac{15}{2}, \frac{31}{2}$$

- “ $\frac{7}{10}$ ” is the Zamolodchikov flow  $\mathcal{M}_4 \rightarrow \mathcal{M}_3$
- “ $\frac{21}{22}$ ” is the coset flow  $[E_8]_2 \rightarrow [E_8]_1$  [Ravanini (1992)]
- “ $\frac{3}{2}$ ” is the massless SShG model which flows from super-Liouville theory (later) [Kim, Rim, Zamolodchikov, CA (2002)]
- New: “ $\frac{31}{2}$ ” is supposed to be a flow from  $E_8$  WZW with level 2; “ $\frac{15}{2}$ ” (?)
- We analyzed other group  $G = su(3), su(4)$  etc to classify all possible UV CFTs based on central charges
- But central charges are **not enough** to identify the UV CFTs

# PART 2. Identifying UV complete QFTs

with Z. Bajnok

# RSOS (non-diagonal) scattering theory

- Consider  $k = p - 2$  with  $G = su(2)$

$$\mathcal{M}_p + \lambda \Phi_{1,3}, \quad \lambda < 0$$

- Particle spectrum: massive kinks

$$a \uparrow b = K_{ab}(\theta), \quad a, b = 0, \frac{1}{2}, \dots, \frac{p}{2} - 1, \quad \text{with } |a - b| = \frac{1}{2}$$

- S-matrix of kinks  $S_{\text{RSOS}}^{[p]}(\theta)$ : non-diagonal [Bernard, LeClair (1990)]

$$K_{ab}(\theta_1) K_{bc}(\theta_2) \rightarrow K_{ad}(\theta_2) K_{dc}(\theta_1)$$

$$S_p(\theta)_{dc}^{ab} = U(\theta) (X_{db}^{ac})^{\frac{i\theta}{2\pi}} \left[ (X_{db}^{ac})^{\frac{1}{2}} \sinh\left(\frac{\theta}{p}\right) \delta_{db} + \sinh\left(\frac{i\pi - \theta}{p}\right) \delta_{ac} \right]$$



# Massless $S$ -matrices [Fendley, Saleur, Zamolodchikov ('93)]

- Massless limit  $\lambda \rightarrow 0^-$

$$S_p^{RR}(\theta) = S_p^{LL}(\theta) = S_p(\theta)$$


- YBE and Crossing-Unitarity relations determine

$$S_p^{RL}(\theta) = S_p^{LR}(-\theta) \propto S_p\left(\theta - \frac{ip\pi}{2}\right)$$

for the  $S$ -matrices of the IR theory  $\mathcal{M}_p + \lambda' \Phi_{3,1}$

- Derived the conjectured TBA (partially) shown before

$$\epsilon_a(\theta) = \frac{\mu R}{2} (\delta_{a0} e^\theta + \delta_{a,p-2} e^{-\theta}) - \sum_{b=0}^{p-2} \mathbb{I}_{ab} \varphi \star \log[1 + e^{-\epsilon_b}](\theta)$$



## TBA for $[T\bar{T}]_s$ deformed minimal CFT $\mathcal{M}_p$





- $[T\bar{T}]_s$  deformations

$$\mathcal{M}_p + \lambda' \Phi_{3,1} + \sum_{s \geq 1} \alpha_s [T\bar{T}]_s$$

- Introduce CDD factors to  $RL, LR$  sectors

$$\tilde{S}_p^{RL}(\theta) = S_{\text{CDD}} \cdot S_p^{RL}(\theta)$$

- Being diagonal,  $S_{\text{CDD}}$  introduces additional kernels between  $R$  &  $L$



## UV complete theories

- Plateaux equations

$$x_n = (1 + x_{n-1})^{1/2}(1 + x_{n+1})^{1/2}, \quad n = 1, \dots, p-3,$$

$$x_0 = (1 + x_1)^{1/2}(1 + x_{p-2})^{\tilde{k}}, \quad x_{p-2} = (1 + x_{p-3})^{1/2}(1 + x_0)^{\tilde{k}}$$

- UV complete only with  $\tilde{k} = \frac{1}{2}$  which means

$$S_{\text{CDD}}^{(1)} = -\tanh\left(\frac{\theta - \beta}{2} - \frac{i\pi}{4}\right) \rightarrow \varphi_{RL} = \frac{1}{\cosh(\theta - \beta)}$$

- We also notice that only real  $\beta$  ( $p > 3$ ) becomes UV complete and

$$c_{\text{eff}} = 3 \frac{p-1}{p+1}$$

## For $p = 3$ , massless Super sinh-Gordon model

- sinh-Gordon model with  $N = 1$  super [Kim, Rim, Zamolodchikov, CA (2002)]

$$\mathcal{L} = \text{Kin.} - \frac{i}{2} \psi \bar{\psi} W''(\phi) + 2\pi [W'(\phi)]^2, \quad W(\phi) = -\mu \sinh(b\phi)$$

- Supersymmetry is spontaneous broken

$$V(\phi) \propto [W'(\phi)]^2 = \cosh^2(b\phi) > 0$$


- massless Goldstino fermion is only spectrum in the IR limit (Ising model)
- TBA is given by

$$\begin{aligned} \epsilon^R(\theta) &= \frac{\mu R}{2} e^\theta - \varphi^{RL} \star \log[1 + e^{-\epsilon^L}], \\ \varphi^{RL}(\theta) &= \frac{1}{\cosh(\theta - ia)} + \frac{1}{\cosh(\theta + ia)}, \quad a = \frac{1 - b^2}{1 + b^2} \end{aligned}$$

Two theories match when  $\beta = 0$  and  $a = 0$



- Ising model  $\mathcal{M}_{p=3}$  deformed by  $[T\bar{T}]_s$  with  $(\beta = 0)$   
= SshG model at self-dual coupling
- Conjecture:  $\mathcal{M}_{p \geq 3}$  deformed by  $[T\bar{T}]_s$  with  $(\beta = 0)$   
=  $\mathbb{Z}_{p-1}$  Parafermionic shG model at self-dual coupling



# Parafermionic shG model

- PshG model


$$\mathcal{L}_{\text{PShG}} = \mathcal{L}_{\text{PF}} + \frac{1}{4\pi}(\partial_\mu\phi)^2 - \kappa \left( \psi_1 \bar{\psi}_1 e^{2b\phi} + \eta \psi_1^\dagger \bar{\psi}_1^\dagger e^{-2b\phi} \right) + \dots$$

- $\psi_1$  is a  $\mathbb{Z}_k$  PF with  $\Delta = 1 - \frac{1}{k}$  ( $k \equiv p - 1$ )
- Fractional SUSY
  - $\eta = 0$  gives Parafermionic LFT [Baseilhac, Fateev (1998)]
  - $\eta = 1$  massive phase
  - $\eta = -1$  massless phase where FSUSY is spontaneously broken
- These types of QFTs compute  $c_{\text{eff}}$  from momentum quantization using Reflection amplitudes

## Reflection amplitudes of LFTs

- Primary fields  $e^{2\alpha\phi}$  and  $e^{2(Q-\alpha)\phi}$  identified upto constant

$$e^{2(\frac{Q}{2}+ip)\phi} = \mathcal{R}(p)e^{2(\frac{Q}{2}-ip)\phi}, \quad \alpha = \frac{Q}{2} + ip, \quad Q = b + \frac{1}{kb}$$




- This amplitude has been computed in the PLFT [Baseilhac, Fateev (1998)]

## Momentum quantization condition

- the perturbation introduces another wall

$$\delta^{(k)}(p) = \pi + 4Qp \ln \frac{R}{2\pi} = \sum_{n=\text{odd}} \delta_n p^n, \quad \mathcal{R}(p) = e^{i\delta^{(k)}(p)}$$



- Scaling function can be obtained from

$$\begin{aligned} c_{\text{eff}} &= \frac{3k}{k+2} - 24p^2 + \mathcal{O}(R) \\ &= \frac{3k}{k+2} - \frac{3\pi^2}{2Q^2 \ln^2 x} - \frac{3\pi^2 \delta_1}{4Q^3 \ln^3 x} - \frac{9\pi^2 \delta_1^2}{32Q^4 \ln^4 x} - \frac{3(2\pi^2 \delta_1^3 + \pi^4 \delta_3)}{64Q^5 \ln^5 x} + \dots \quad (x = \frac{R}{2\pi}) \end{aligned}$$



## Reflection amplitude vs. massless TBA

- Need to solve numerically with high precision


$$c_{\text{eff}}(r) = \frac{3k}{k+2} + \sum_{n=2} \frac{c_n(k)}{(\log r)^n} + O(r)$$

- Two match very well at self-dual coupling  $b = \frac{1}{\sqrt{k}}$


	$c_2$	$c_3$	$c_4$	$c_5$
$k = 2$	7.402199	42.7620	185.218	714.563
	7.402203	42.7628	185.282	717.247
$k = 3$	11.10332	77.8573	409.598	1924.84
	11.10330	77.8543	409.425	1919.36
$k = 4$	14.8045	119.428	722.797	3898.75
	14.8044	119.4197	722.475	3892.55

## What if $\beta \neq 0$ for generic $p \geq 3$ ?

- Numerical solutions of the massless TBA show “Roaming” [ $p = 4$ ,  $\beta = 10\pi$ ]



## Roaming TBAs with $\beta \gg 1$



- Each plateau corresponds to

$$\mathcal{M}_{k+1} \rightarrow \mathbb{Z}_k \mathcal{M}_1 \rightarrow \mathbb{Z}_k \mathcal{M}_{k+1} \rightarrow \mathbb{Z}_k \mathcal{M}_{2k+1}$$

where

$$\mathbb{Z}_k \mathcal{M}_\ell = \frac{su(2)_k \otimes su(2)_\ell}{su(2)_{k+\ell}}$$

- Very similar with PF staircase TBA by [Dorey, Ravanini (1993)]
- A difference occurs at the first plateau (IR CFT) which is a minimal CFT

## Discussions

- We have found exact massless  $S$ -matrices from which we have derived TBA for new RG flows
- We have found that there is only one new RG flow to  $\mathcal{M}_p$  for  $p > 3$  if the CFT is in RSOS (can not exclude existence of other basis)
- and we have identified those as PF sinh-Gordon ( $\beta = 0$ ) and PF Roaming ( $\beta \gg 1$ ) models
- Can we find exact  $S$ -matrices in this way for all conjectured TBA and NLIE?
- Can we generalize this approach to other irrelevant deformations than  $T\bar{T}$ 's?

Thank you for attention!